

LITERATURE FORMING THE BASIS FOR *WHERE RIVERS ARE BORN*

- Adams, S.B., and C.A. Frissell. 2001. Thermal habitat use and evidence of seasonal migration by Rocky Mountain tailed frogs, *Ascaphus montanus*, in Montana. *Canadian Field-Naturalist* 115: 251-256.
- Adams, S.B., C.A. Frissell, and B.E. Rieman. 2001. Geography of invasion in mountain streams: consequences of headwater lake fish introductions. *Ecosystems* 4: 296-307.
- Alexander, D.R., and H.R. MacCrimmon. 1974. Production and movement of juvenile rainbow trout (*Salmo gairdneri*) in a headwater of Bothwell's Creek, Georgian Bay, Canada. *Journal of the Fisheries Research Board of Canada* 31: 117-121.
- Amon, J.P., C.A. Thompson, Q.J. Carpenter, and J. Miner. 2002. Temperate zone fens of the glaciated Midwestern USA. *Wetlands* 22(2): 301-317.
- Arnold, C. L., P.J. Boison, and P.C. Patton. 1982. Sawmill Brook – An example of rapid geomorphic change related to urbanization. *Journal of Geology* 90:155-166.
- Bärlocher, F., and M.A.S. Graca. 2002. Exotic riparian vegetation lowers fungal diversity but not leaf decomposition in Portuguese streams. *Freshwater Biology* 47: 1123-1135.
- Bedford, B.L, M.R. Walbridge, and A. Aldous. 1999. Patterns in nutrient availability and plant diversity of temperate North American wetlands. *Ecology* 80(7): 2151-2169.
- Bedford, B.L., D.J. Leopold, and J.P Gibbs. 2001. Wetland ecosystems. P. 781-804. *In* Encyclopedia of Biodiversity, Volume 5. Academic Press, San Diego, CA, USA.
- Beer, T. and M. Borgas. 1993. Horton laws and the fractal nature of streams. *Water Resources Research* 29:1475-1487.
- Belk, D. 1996. Was sind "Urzeitkrebse"? *Stapfia* 42: 15-19.
- Berven, K.A. and T.A. Grudzien. 1990. Dispersal in the wood frog (*Rana sylvatica*): Implications for genetic population structure. *Evolution* 44(8): 2054-2056.
- Biek, R., L.S. Mills, and R.B. Bury. 2002. Terrestrial and stream amphibians across clearcut forest interfaces in the Siskiyou Mountains, Oregon. *Northwest Science* 76: 129-140.
- Bilby, R. E. and G. E. Likens. 1980. Importance of organic debris dams in the structure and function of stream ecosystems. *Ecology* 61:1107-1113.
- Bilby, R. E. 1981. Role of organic debris dams in regulating the export of dissolved and particulate matter from a forested watershed. *Ecology* 62:1234-1243.

- Bilby, R. E., and G. E. Likens. 1979. Effect of hydrologic fluctuations on the transport of fine particulate organic carbon in a small stream. *Limnol. Oceanogr.* 24:69-75.
- Blaustein, A.R., D.B. Wake, and W.P. Sousa. 1994. Amphibian declines—judging stability persistence, and susceptibility of populations to local and global extinctions. *Conservation Biology* 8(1): 60-71.
- Bott, T. L., L. A. Kaplan, and F. T. Kuserk. 1984. Benthic bacterial biomass supported by streamwater dissolved organic matter. *Microbial Ecology* 10:335-344.
- Bott, T. L., J. T. Brock, C. S. Dunn, R. J. Naiman, R. W. Ovink, and R. C. Petersen. 1985. Benthic community metabolism in four temperate stream systems: An inter-biome comparison and evaluation of the river continuum concept. *Hydrobiologia* 123:3-45.
- Brown, K.S. 1998. Vanishing pools taking species with them. *Science* 281:626
- Buhlmann, K.A., and J.W. Gibbons. 1997. Imperiled aquatic reptiles of the southeastern United States: historical review and current conservation status. Pages 201-232 *in*: G.W. Benz and D.E. Collins (editors). *Aquatic fauna in peril: the southeastern perspective*. Southeast Aquatic Research Institute Special Publication 1. Lenz Design and Communications, Decatur, GA.
- Burton, T.M. and G.E. Likens. 1975. Salamander populations and biomass in the Hubbard Brook Experimental Forest New Hampshire, USA. *Copeia* (3): 541-546.
- Bury, R.B. 1988. Habitat relationships and ecological importance of amphibians and reptiles. Pages 61-76 *in*: K.J. Raedeke (editor). *Streamside management: riparian wildlife and forestry interactions*. University of Washington Institute of Forest Resources Contribution 59.
- Carpenter, Q. 1995. Toward a new definition of calcareous fen for Wisconsin (USA). Ph.D. Dissertation. Gaylord Nelson Institute for Environmental Studies, University of Wisconsin, Madison, WI, USA.
- Carter, B., and W.A. Hubert. 1995. Factors influencing fish assemblages of a high-elevation desert stream system in Wyoming. *Great Basin Naturalist* 55: 169-173.
- Cedfeldt, P.T., M.C. Watzin, and B.D. Richardson. 2000. Using GIS to identify functionally significant wetlands in the Northeastern United States *Environmental Management* 26 (1): 13-24.
- Cherry and Freeze 1979. *Groundwater*.
- Chou, R.Y.M., L.C. Ferrington, B.L. Hayford, and H.M. Smith. 1999. Composition and phenology of Chironomidae (Diptera) from an intermittent stream in Kansas. *Archiv für Hydrobiologie* 147: 35-64.

- Coffman, D. M., E. A. Keller, and W. N. Melhorn. 1972. New topologic relationship as an indicator of drainage network evolution. *Water Resources Research* 8:1497-1505.
- Coffman, W.P., and C.L. de la Rosa. 1998. Taxonomic composition and temporal organization of tropical and temperate species assemblages of lotic Chironomidae. *Journal of the Kansas Entomological Society* 71: 388-406.
- Cole, J. J. and N. F. Caraco. 2001. Carbon in catchments: connecting terrestrial carbon losses with aquatic metabolism. *Marine and Freshwater Research*. 52:101-110.
- Cross, S.P. 1988. Riparian systems and small mammals and bats. Pages 93-112 *in*: K.J. Raedeke (editor). *Streamside management: riparian wildlife and forestry interactions*. University of Washington Institute of Forest Resources Contribution 59.
- Curry, R.A., C. Brady, D.L.G. Noakes, and R.G. Danzmann. 1997. Use of small streams by young brook trout spawned in a lake. *Transactions of the American Fisheries Society* 126: 77-83.
- Cushing, C.E. 1996. The ecology of cold desert spring-streams. *Archiv für Hydrobiologie* 135: 499-522.
- Delucchi, C.M. 1988. Comparison of community structure across streams with different temporal flow regimes. *Canadian Journal of Zoology* 66: 579-586.
- Department of Interior. 1968. *The Nation's River*. The Department of Interior Official Report on the Potomac. O – 310 – 865. Washington, D.C.: U. S. Government Printing Office.
- Dieterich, M. Anderson N. H. 1998. Dynamics of abiotic parameters, solute removal and sediment retention in summer-dry headwater streams of western Oregon. *Hydrobiologia* 379:1-15.
- Dieterich, M. Anderson N. H. 2000. The invertebrate fauna of summer-dry streams in western Oregon. *Archiv für Hydrobiologie* 147:273-295.
- Dietrich, W. E. and Dunne T. 1978. Sediment budget for small catchment in mountainous terrain. *Zeitschrift für Geomorphologie Neue Folge, Supplement band* 29:191-206.
- Dietrich, W. E. and Dunne T. 1993. The channel head. Pages 176-219 *in* K. Beven and M. J. Kirkby, editors. *Channel network hydrology*. New York: John Wiley and Sons.
- Dobson, M., A. G. Hildrew, A. Ibbotson, and J. Garthwaite. 1992. Enhancing litter retention in streams: do altered hydraulics and habitat area confound field experiments. *Freshwater Biology* 28:71-79.

- Dobson, M., and A. G. Hildrew. 1992. A test of resource limitation among shredding detritivores in low order streams in Southern England. *Journal of Animal Ecology*, 61:69-77.
- Dodd, C.K. 1997. Imperiled amphibians: a historical perspective. Pages 165-200 *in*: G.W. Benz and D.E. Collins (editors). *Aquatic fauna in peril: the southeastern perspective*. Southeast Aquatic Research Institute Special Publication 1. Lenz Design and Communications, Decatur, GA.
- Dodd, C.K. Jr. 1992. Biological diversity of a temporary pond herpetofauna in north Florida sandhills. *Biodiversity and Conservation* 1:125-142.
- Dodd, C.K. Jr. and B.S. Cade. 1998. Movement patterns and the conservation of amphibians breeding in small temporary wetlands. *Conservation Biology* 12(2): 331-339.
- Drexler, J.Z. and B.L. Bedford. 2002. Pathways of nutrient loading and impacts on plant diversity in a New York peatland. *Wetlands* 22(2): 263-281.
- Dubey, T., S.L. Stephenson, and P.J. Edwards. 1994. Effect of pH on the distribution and occurrence of aquatic fungi in six West Virginia mountain streams. *Journal of Environmental Quality* 23: 1271-1279.
- Ducks Unlimited, Inc. 2001. http://www.ducks.org/conservation/404_report.asp.
- Duijsings, J. J. H. M. 1987. A sediment budget for a forested catchment and its implications for channel development. *Earth Surfaces Processes and Landforms* 12:173-184.
- Dunne, T. 1991. Stochastic aspect of the relations between climate, hydrology, and landform evolution. *Transactions of the Japanese Geomorphological Union* 12:1-24.
- Dunne, T. and L. B. Leopold. 1978. *Water in environmental planning*. New York: W. H. Freeman and Company.
- Dunstone, N., and M.L. Gorman (editors). *Behaviour and ecology of riparian mammals*. Cambridge University Press, UK. 391 pp.
- Edwards, R. T. 1998. The hyporheic zone. Pages 399-429 *in* R. J. Naiman and R. E. Bilby, editors. *River ecology and management: Lessons from the Pacific coastal ecoregion*. New York: Springer Verlag.
- Elwood, J. W., J. D. Newbold, A. F. Trimble, and R. W. Stark. 1981. The limiting role of phosphorus in a woodland stream ecosystem: Effects of P enrichment on leaf decomposition and primary producers. *Ecology* 62:146-158.

- Erman, D.C., and G.R. Leidy. 1975. Downstream movement of rainbow trout fry in a tributary of Sagehen Creek under permanent and intermittent flow. *Transactions of the American Fisheries Society* 104: 467-473.
- Erman, D.C., and V.M. Hawthorne. 1976. The quantitative importance of an intermittent stream in the spawning of rainbow trout. *Transactions of the American Fisheries Society* 105: 675-681.
- Erman, N.A., and D.C. Erman. 1995. Spring permanence, Trichoptera species richness, and the role of drought. *Journal of the Kansas Entomological Society* 68: 50-64.
- Etnier, D.A. 1997. Jeopardized southeastern freshwater fishes: a search for causes. Pages 87-104 *in*: G.W. Benz and D.E. Collins (editors). *Aquatic fauna in peril: the southeastern perspective*. Southeast Aquatic Research Institute Special Publication 1. Lenz Design and Communications, Decatur, GA.
- Fausch, K.D., and R.G. Bramblett. 1991. Disturbance and fish communities in intermittent tributaries of a western Great Plains river. *Copeia* 1991: 659-674.
- Feminella, J.W. 1996. Comparison of benthic macroinvertebrate assemblages in small streams along a gradient of flow permanence. *Journal of the North American Benthological Society* 15: 651-669.
- Fend, S.V., and D.L. Gustafson. 2001. A new lumbriculid genus and species from North America (Clitellata, Lumbriculidae). *Hydrobiologia* 463: 13-22.
- Fend, S.V., and R.O. Brinkhurst. 2000. New species of *Rhynchelmis* (Clitellata, Lumbriculidae), with observations on the Nearctic species. *Hydrobiologia* 428: 1-59.
- Fetherston, K. L., R. J. Naiman, and R. E. Bilby. 1995. Large woody debris, physical process, and riparian forest development in montane river networks of the Pacific Northwest. *Geomorphology* 13:133-144.
- Fisher, S. G. 1977. Organic matter processing by a stream-segment ecosystem: Fort River, Massachusetts, U. S. A. *Int. Revue ges. Hydrobiol.* 62:701-727.
- Fisher, S. G. 1997. Creativity, idea generation, and the functional morphology of streams. *Journal of the North American Benthological Society* 16:305-318.
- Fisher, S. G., and G. E. Likens. 1973. Energy flow in Bear Brook, New Hampshire: An integrative approach to stream ecosystem metabolism. *Ecol. Monogr.* 43:421-439.
- Flinders, C.A., and D.D. Magoulick. 2003. Effects of habitat permanence on crayfish community structure. *American Midland Naturalist* 149: 134-147.

- Fritz, K.M., and W.K. Dodds. 2002. Macroinvertebrate assemblage structure across a tallgrass prairie stream landscape. *Archiv für Hydrobiologie* 154: 79-102.
- Gary, M., R. McAfee, Jr. and C. L. Wolf, editors. 1974. *Glossary of Geology*. Washington, DC: American Geological Institute.
- Gibbons, J.W. and D.H. Bennett. 1974. Determination of anuran terrestrial activity patterns by a drift fence method. *Copeia* (1): 236-243.
- Gibbons, J.W. and R.D. Semlitsch. 1981. Terrestrial drift fences with pitfall traps : An effective technique for quantitative sampling of animal populations. *Brimleyana* 7:1-16.
- Gibbs, J.P. 1993. Importance of small wetlands for the persistence of local populations of wetland-associated animals. *Wetlands*. 13:25-31.
- Glazier, D.S. 1991. The fauna of North American temperate cold springs: patterns and hypotheses. *Freshwater Biology* 26: 527-542.
- Godwin, K.S., J.P. Shallenberger, D.J. Leopold, and B.L. Bedford. 2002. Linking landscape properties to local hydrogeologic gradients and plant species occurrence in minerotrophic fens of New York State, USA: a hydrogeologic setting (HGS) framework. *Wetlands* 24: 722-737.
- Gomi, T., R. C. Sidle, and J. S. Richardson. 2002. Understanding processes and downstream linkages of headwaters systems. *BioScience* 52:905-916.
- Gomi, T., R. C. Sidle, M. D. Bryant, and R. D. Woodsmith. 2001. The characteristics of woody debris and sediment distribution in headwater streams, south-eastern Alaska. *Canadian Journal of Forest Research* 31:1386-1399.
- Gray, L.J. 1993. Response of insectivorous birds to emerging aquatic insects in riparian habitats of a tallgrass prairie stream. *American Midland Naturalist* 129: 288-300.
- Hack, J. T. and J. C. Goodlett. 1960. *Geomorphology and Forest Ecology of a Mountain Region in the Central Appalachians*. Washington, D. C.: U. S. Geological Survey, Professional paper no. 347.
- Haggerty, S.M., D.P. Batzer, and C.R. Jackson. 2002. Macroinvertebrate assemblages in perennial headwater streams of the Coastal Mountain range of Washington, USA. *Hydrobiologia* 479: 143-154.
- Hall, R.O., G.E. Likens, and H.M. Malcom. 2001. Trophic basis of invertebrate production in 2 streams at the Hubbard Brook Experimental Forest. *Journal of the North American Benthological Society* 20: 432-447.
- Hansen, W. F. 2001. Identifying stream types and management implications. *Forest Ecology and Management* 143:39-46.

- Harvey, M.J., and J.D. Clark. 1997. Imperiled mammalian fauna of aquatic ecosystems in the Southeast: a historical perspective. Pages 245-258 *in*: G.W. Benz and D.E. Collins (editors). Aquatic fauna in peril: the southeastern perspective. Southeast Aquatic Research Institute Special Publication 1. Lenz Design and Communications, Decatur, GA.
- Hedges, J. I., H. R. Ertel, P. D. Quay, P. M. Grootes, J. E. Richey, A. H. Devol, G. W. Farwell, F. W. Schmidt, and E. Salati. 1986. Organic carbon-14 in the Amazon River system. *Science* 231:1129-31.
- Helmlinger, K. R., P. Kumar, and E. Foufoulageorgiou. 1993. On the use of digital elevation model data for Hortonian and fractal analyses of channel networks. *Water Resources Research* 29:2599-2613.
- Herschler, R. 1994. A review of the North American freshwater snail genus *Pyrgulopsis* (Hydrobiidae). *Smithsonian Contributions to Zoology* 554: 1-115.
- Hey, R. D. 1979. Dynamic process-response model of river channel development. *Earth Surface Processes* 4:59-72.
- Hickman, J.C., Editor. 1993. The Jepson manual of higher plants of California. University of California Press, Berkeley, CA, USA.
- Hill, B.H., R.J. Stevenson, Y. Pan, A.T. Herlihy, P.R. Kaufmann, and C.B. Johnson. 2001. Comparison of correlations between environmental characteristics and stream diatom assemblages characterized at genus and species levels. *Journal of the North American Benthological Society* 20: 299-310.
- Horton, R. E. 1945. Erosional development of streams and their drainage basins: hydrophysical approach to quantitative morphology. *Geological Society of America Bulletin* 56:275-370.
- Jass, J., and B. Klausmeier. 2000. *Atlas and bibliography of the first state and county records for Anostracans (Crustacea: Branchiopoda) of the contiguous United States. Contributions In Biology and Geology No. 94.* Milwaukee County Museum, Milwaukee, WI, USA.
- Johnson, P.D., and K.M. Brown. 2000. The importance of microhabitat factors and habitat stability to the threatened Louisiana pearl shell, *Margaritifera hembeli* (Conrad). *Canadian Journal of Zoology* 78: 271-277.
- Kaplan, L.A., and J. D. Newbold. 2002. The Role of Monomers in Stream Ecosystem Metabolism, pp. 97-119, In: Aquatic Ecosystems Interactivity of Dissolved Organic Matter, S.E.G. Findlay and R. L. Sinsabaugh, ed., Academic Press.
- Kaplan, L.A., R. A. Larson, and T. L. Bott. 1980. Patterns of dissolved organic carbon in transport. *Limnology and Oceanography* 25:1034-1043.

- Kaushik, N. K., and H. B. N. Hynes. 1971. The fate of the dead leaves that fall into streams. *Archiv fur Hydrobiologie* 68:465-515.
- Keeler-Wolf, T., D.R. Elam, K.Lewis, and S.A. Flint. 1998. *California vernal pool assessment: Preliminary report*. The Resources Agency, Department of Fish and Game, Sacramento, CA, USA.
- Kenney, L.P. and M.R. Burne. 2000. A Field Guide to Vernal Pools. Massachusetts Division of Fisheries and Wildlife and Natural Heritage and Endangered Species Program. Westborough, Massachusetts.
- Kiffney, P. M., J. S. Richardson, and M. C. Feller. 2000. Fluvial and epilithic organic matter dynamics in headwater streams of southwestern British Columbia, Canada. *Arch. Hydrobiol.* 149:109-129.
- Kirkby M. J. 1993. Network hydrology and geomorphology. Pages 1-11 *in* K. Beven and M. J. Kirkby, editors. *Channel Network Hydrology*. New York: John Wiley and Sons.
- Kirkman, L.K., P.C. Goebel, L. West, M.B. Drew, and B.J. Palik. 2000. Depressional wetland vegetation types: A question of plant community development. *Wetlands* 20(2):373-385.
- Knox, J. C. 1977. Human impacts on Wisconsin stream channels. *Annals of the Association of American Geographers* 67:323-342.
- Knox, J. C., P. J. Bartlein, and W. C. Johnson. 1974. Environmental assessment of sediment sources and sedimentation distributions for the Lake LaFarge watershed and impoundment; in *Environmental Analysis of the Kickapoo River Impoundment*: Institute for Environmental Studies, Univ. of Wisconsin, Report 28, pp. 77-116.
- Kondratieff, B.C. and C.J. Pyott. 1987. The Anisoptera of the Savannah River Plant, South Carolina, USA: Thirty years later. *Odonatologica* 16: 9-24.
- Krause, C.G., W.A. Hubert, and F.J. Rahel. 2001. An assessment of headwater isolation as a conservation strategy for cutthroat trout in the Absaroka Mountains of Wyoming. *Northwest Science* 75: 1-11.
- Krusic, R.A., M. Yamasaki, C.D. Neefus, and P.J. Pekins. 1996. Bat habitat use in White Mountain National Forest. *Journal of Wildlife Management* 60: 625-631.
- Kruuk, H., D.N. Carss, J.W.H. Conroy, and M.J. Gaywood. 1998. Habitat use and conservation of otters (*Lutra lutra*) in Britain: a review. Pages 119-133 *in*: N. Dunstone and M.L. Gorman (editors). *Behaviour and ecology of riparian mammals*. Cambridge University Press, UK. 391 pp.
- Kuichling, E. 1889. The relationship between the rainfall and discharge of sewers in populous districts. *Transactions of the American Society of Civil Engineers*, v. 20.

- Labbe, T.R., and K.D. Fausch. 2000. Dynamics of intermittent stream habitat regulate persistence of a threatened fish at multiple scales. *Ecological Applications* 10: 1774-1791.
- Lamberti, G.A. 1996. The role of periphyton in benthic food webs. Pages 533-572 *in*: R.J. Stevenson, M.L. Bothwell, and R.L. Lowe (editors). *Algal ecology: freshwater benthic ecosystems*. Academic Press.
- Leehner, J.A., R. L. Malcolm, P. W. McKinley, and L. A. Eccles. 1974. Occurrence of dissolved organic carbon in selected groundwater samples in the United States. *J. Res. U.S. Geol. Survey* 2:361-369.
- Leopold, L. B. 1968. *Hydrology for urban land planning—a guidebook on the hydrologic effects of urban land use*. U. S. Geologic Survey Circular 554, Washington, D. C.: U. S. Government Printing Office.
- Leopold, L. B. 1994. *A view of the river*. Cambridge, Mass: Harvard University Press.
- Leopold, L. B. and T. Maddock, Jr. 1953. *The hydraulic geometry of stream channels and some physiographic implications*. U.S. Geological Survey Professional Paper 252, Washington, D.C.: U. S. Government Printing Office.
- Leopold, L. B., M. G. Wolman, and J. P. Miller. 1964. *Fluvial processes in geomorphology*. San Francisco: W. H. Freeman.
- Lotrich, V.A. 1973. Growth, production, and community composition of fishes inhabiting a first-, second-, and third-order stream of eastern Kentucky. *Ecological Monographs* 43: 377-397.
- Lovett, G. M., K. C. Weathers, and W. Sobczak. 2000. Nitrogen saturation and retention in forested watersheds of the Catskill Mountains, New York. *Ecological Applications* 10:73-84.
- Lowe, W.H., and D.T. Bolger. 2002. Local and landscape-scale predictors of salamander abundance in New Hampshire headwater streams. *Conservation Biology* 16: 183-193.
- Lowrance, R.R., R.L. Todd, J. Fair, Ol Hendrickson, R. Leonard and L. Asmussen. 1984. Riparian forests as nutrient filters in agricultural watersheds. *BioSciences* 34: 374-377.
- Lowrance, R.R. and others. 1997. Water quality functions of riparian forest buffers in Chesapeake Bay watersheds. *Environmental Management* 21: 687 - 712.
- Mahoney, D.L., D.A. Mort, and B.E. Taylor. 1989. Species richness of calanoid copepods, cladocerans, and other branchiopods in Carolina bay temporary ponds. *American Midland Naturalist* 123:244-258.

- Marani, A., R. Rigon, and A. Rinaldo. 1991. A note on fractal channel networks. *Water Resources Research* 27:3041-3049.
- Master, L.L., B.A. Stein, L.S. Kutner, and G.A. Hammerson. 2000. Vanishing assets: conservation status of U.S. species. Pages 93-118 *in*: B.A. Stein, L.S. Kutner, and J.S. Adams (editors). *Precious heritage: the status of biodiversity in the United States*. Oxford University Press.
- Matthews, W.J. 1998. *Patterns in freshwater fish ecology*. Chapman and Hall.
- Matthews, W.J., and J.D. Maness. 1979. Critical thermal maxima, oxygen tolerances and success of cyprinid fish in a southwestern river. *American Midland Naturalist* 102: 374-377.
- Matthews, W.J., and J.T. Styron. 1981. Tolerance of headwater vs. mainstream fishes for abrupt physicochemical changes. *American Midland Naturalist* 105: 149-158.
- Mayer, M.S., and G.E. Likens. 1987. The importance of algae in a shaded headwater stream as food for an abundant caddisfly. *Journal of the North American Benthological Society* 6: 262-269.
- McCabe, D.J., and J.L. Sykora. 2000. Community structure of caddisflies along a temperate springbrook. *Archiv für Hydrobiologie* 148: 263-282.
- McKey-Fender, D., and W.M. Fender. 1988. *Phagodrilus* gen. nov. (Lumbriculidae) – systematics and biology of a predaceous oligochaete from western North America. *Canadian Journal of Zoology* 66: 2304-2311.
- Megahan, W. F. 1982. Channel sediment storage behind obstructions in forested drainage basins draining the granitic bedrock of the Idaho Batholith. Pages 114-121 *in* F. J. Swanson, R. J. Janda, T. Dunne, and D. N. Swanston, editors. *Sediment budgets and routing in forested drainage basins*. US Department of Agriculture Forest Service general technical report PNW-141.
- Meinzer, O. E. 1923. *Outline of groundwater hydrology, with definitions*. U.S. Geological Survey. Water-supply Paper 494. Washington, D. C.: U. S. Government Printing Office.
- Meyer, J. L. and J. B. Wallace. 2001. Lost linkages and lotic ecology: rediscovering small streams. Pages 295-317 *in* M.C. Press, N.J. Huntly, and S. Levin, editors. *Ecology: achievement and challenge*. Blackwell Science.
- Minckley, W.L. 1999. Ecological review and management recommendations for recovery of the endangered Gila topminnow. *Great Basin Naturalist* 59: 230-244.
- Minshall, G. W. 1978. Autotrophy in stream ecosystems. *Bioscience* 28:767-771.

- Moler, P.E., and R. Franz. 1987. Wildlife values of small, isolated wetlands in the southeastern coastal plain. Pp. 234-241 in R. Odum, K.A. Riddleberger, and J.C. Ozier, eds. Proc. Third Southeastern Nongame and Endangered Wildlife Symposium. Georgia Dept. of Natural Resources, Atlanta.
- Montgomery, D. R. 1999. Process domain and river continuum. *Journal of the American Water Resources Association* 35:397-410.
- Montgomery, D. R. and E. Foufoula-Georgiou. 1993. Channel network source representation using digital elevation models. *Water Resources Research* 29:3925-3934.
- Montgomery, D. R. and W. E. Dietrich. 1989. Source areas, drainage density, and channel initiation. *Water Resources Research* 25:1907-1918.
- Montgomery, D., and W. E. Dietrich. 1992. Channel initiation and the problem of landscape scale. *Science* 255:826-830.
- Moore, D.R.J., P. Keddy, C. Gaudet and I. Wisheu. 1989. Conservation of wetlands: Do infertile wetlands deserve a higher priority? *Biological Conservation* 47:203-217.
- Morisawa, M. E. 1957. Accuracy of determination of stream lengths from topographic maps. *Eos, Transactions of the American Geophysical Union* 38: 86-88.
- Morse, J.C., B.P. Stark, W.P. McCafferty, and K.J. Tennseen. 1997. Southern Appalachian and other southeastern streams at risk: implications for mayflies, dragonflies and damselflies, stoneflies, and caddisflies. Pages 17-42 in: G.W. Benz and D.E. Collins (editors). *Aquatic fauna in peril: the southeastern perspective*. Southeast Aquatic Research Institute Special Publication 1. Lenz Design and Communications, Decatur, GA.
- Murray, N.L., and D.F. Stauffer. 1995. Nongame bird use of habitat in central Appalachian riparian forests. *Journal of Wildlife Management* 59: 78-88.
- Naiman, R. J. 1976. Primary production, standing stock, and export of organic matter in a Mohave Desert thermal stream. *Limnol. Oceanogr.* 21:60-73.
- NatureServe. 2003. www.natureserve.org [accessed February 2003].
- Nekola, J.C. 1994. The environment and vascular flora of northeastern Iowa fen communities. *Rhodora* 96(886): 121-169.
- Newbold, J. D. 1992. Cycles and spirals of nutrients. pp. 379-408 in P. Calow and G. E. Petts, editors. *The rivers handbook*. Volume 1. Hydrological and ecological principles. Blackwell Scientific, Oxford.
- Newbold, J. D., P. J. Mulholland, J. W. Elwood, and R. V. O'Neill. 1982. Organic carbon spiralling in stream ecosystems. *Oikos* 38:266-272.

- Niyogi, D.K., D.M. McKnight, and W.M. Lewis. 2002. Fungal communities and biomass in mountain streams affected by mine drainage. *Archiv für Hydrobiologie* 155: 255-271.
- Nussbaum, R.A. 1977. Aspects of life history and ecology of Olympic salamander, *Rhyacotriton olympicus* (Gage). *American Midland Naturalist* 98: 176-199.
- Ohio Environmental Protection Agency. 2002a. Clean rivers spring from their source: the importance and management of headwaters streams. Columbus, Ohio: State of Ohio Environmental Protection Agency, Division of Surface Water.
- Ohio Environmental Protection Agency. 2002b. The importance and benefits of primary headwater streams. Columbus, Ohio: State of Ohio Environmental Protection Agency, Division of Surface Water.
- Ohio Environmental Protection Agency. 2002c. Nonpoint source impacts on primary headwater streams. Columbus, Ohio: State of Ohio Environmental Protection Agency, Division of Surface Water.
- Ohio Environmental Protection Agency. 2002d. Ohio EPA primary headwater habitat initiative data compendium 1999-2000: habitat, chemistry, and stream morphology data. Columbus, Ohio: State of Ohio Environmental Protection Agency, Division of Surface Water.
- Ostrand, K.G., and G.R. Wilde. 2001. Temperature, dissolved oxygen, and salinity tolerances of five prairie stream fishes and their role in explaining fish assemblage patterns. *Transactions of the American Fisheries Society* 130: 742-749.
- Paller, M.H. 1994. Relationships between fish assemblage structure and stream order in South Carolina Coastal Plain streams. *Transactions of the American Fisheries Society* 123: 150-161.
- Pavelis, G. A. 1987. Economic survey of farm drainage. Pages 110-136 in G. A. Pavelis, editor. *Farm drainage in the United States: history, status, and prospects*. Washington, D. C.: U. S. Dept. of Agriculture, Misc. Pub. 1455.
- Perry, C. A., D. M. Wolock and J. C. Artman. 2002. Estimates of median flows for streams on the Kansas Surface Water Register. U.S. Geological Survey Water-Resources Investigation Report 02-4292, Washington, D.C.: U.S. Government Printing Office.
- Peterjohn, W.T. and D.L. Correll. 1984. Nutrient dynamics in an agricultural watershed: Observation on the role of a riparian forest. *Ecology* 65: 1466 - 1475.
- Petts, G. E. and C. Amoros. 1996. The fluvial hydrosystem. Pages 1-12 in G. E. Petts, and C. Amoros, editors. *Fluvial Hydrosystems*. London: Chapman and Hall.

- Pigg, K. B. 1992. Evolution of Isoetalian lycopsids. *Annals of the Missouri Botanical Garden* 79: 789-612.
- Poiani, K.A., B.D. Richter, and M.G. Anderson. 2000. Biodiversity conservation at multiple scales: Functional sites, landscapes, and networks. *BioScience* 50: 133-146.
- Pounds, J.A. and M.L. Crump. 1994. Amphibian declines and climate disturbance—the case of the golden toad and the harlequin frog. *Conservation Biology* 8 (1): 72-85.
- Progar, R.A., and A.R. Moldenke. 2002. Insect production from temporary and perennially flowing headwater streams in western Oregon. *Journal of Freshwater Ecology* 17: 391-407.
- Raedeke, K.J., R.D. Taber, and D.K. Paige. 1988. Ecology of large mammals in riparian systems of Pacific Northwest forests. Pages 113-132 *in*: K.J. Raedeke (editor). *Streamside management: riparian wildlife and forestry interactions*. University of Washington Institute of Forest Resources Contribution 59.
- Raymond, P. A., and J. E Bauer. 2001. Use of ^{14}C and ^{13}C natural abundances for evaluating riverine, estuarine, and coastal DOC and POC sources and cycling: a review and synthesis. *Organic Geochemistry* 32:469-485.
- Reice, S. R. 1978. Role of detritivore selectivity in species-specific litter decomposition in a woodland stream. *Verhandlungen Internationale Vereinigung für Theoretische und Angewandte Limnologie*. 20:1396-1400.
- Richards, K. 1993. Sediment delivery and the drainage network. Pages 221-254 in K. Beven and M. J. Kirkby, editors. *Channel Network Hydrology*. New York: John Wiley and Sons.
- Richardson, J. S. 1992. Food, microhabitat, or both? Macroinvertebrate use of leaf accumulations in a montane stream. *Freshwater Biology* 27:169-176.
- Riley, A. L. 1998. *Restoring streams in cities: A guide for planners, policymakers, and citizens*. Washington, D.C.: Island Press.
- Roni, P. 2002. Habitat use by fishes and Pacific giant salamanders in small western Oregon and Washington streams. *Transactions of the American Fisheries Society* 131: 743-761.
- Russell, K.R., D.C. Guynn, and H.G. Hanlin. 2002. The importance of small isolated wetlands for herpetofaunal diversity in managed, young growth forests in the Coastal Plain of South Carolina. *Forest Ecology and Management* 163(1-3): 43-59.
- Ryon, M.G. 1986. The life history and ecology of *Etheostoma trisella* (Pisces: Percidae). *American Midland Naturalist* 115: 73-86.
- Sanzone, D.M., J.L. Meyer, E. Marti, E.P. Gardiner, J.L. Tank, and N.B. Grimm. 2003. Carbon and nitrogen transfer from a desert stream to riparian predators. *Oecologia* 134: 238 - 250.

- Scott, A. F., and A. Bufalino. 1997. Dynamics of the amphibian communities at two small ponds in Land Between the Lakes over the past decade. *In* A. F. Scott, S.W. Hamilton, E.W. Chester, and D.S. White, eds. Proc. Seventh Symposium on the Natural History of Lower Tennessee and Cumberland River valleys. Austin Peay State University, Clarksville, Tennessee.
- Seidman, V.M., and C.J. Zabel. 2001. Bat activity along intermittent streams in northwestern California. *Journal of Mammalogy* 82: 738-747.
- Semlitsch, R.D. 1997. Biological delineation of terrestrial buffer zones for pond breeding salamanders. *Conservation Biology* 12(4): 1113-1119.
- Semlitsch, R.D. 2000. Size does matter: The value of small isolated wetlands. *National Wetlands Newsletter* January-February 5-6.
- Semlitsch, R.D., and J.R. Bodie. 1998. Are small, isolated wetlands expendable? *Conservation Biology* 12:1129-1133.
- Shafroth, P.B., J.C. Stromberg and D.T. Patten. 2002. Riparian vegetation response to altered disturbance and stress regimes. *Ecological Applications* 12: 107 - 123.
- Sharitz, R.R., and J.W. Gibbons. 1982. The ecology of southeastern shrub bogs (pocosins) and Carolina bays: A community profile. FWS/OBS-82/04. U.S. Fish and Wildlife Service, Washington, D.C.
- Shimano, Y. 1992. Characteristics of the Stream Network Composition of Drainage Basins in the Japanese Islands. *Environmental Geology and Water Sciences* 20:5-14.
- Shreve, R. L. 1967. Infinite topologically random channel networks. *Journal of Geology* 75:178-186.
- Sidle, R. C, Y. Tsuboyama, S. Noguchi, I. Hosoda, M. Jujieda, and T. Shimizu. 2000. Streamflow generation in steep headwaters: A linked hydro-geomorphic paradigm. *Hydrological Processes* 14:369-385.
- Soil Conservation Service, U. S. Department of Agriculture. 1986. Urban hydrology for small watersheds. USDA-SCS Engineering Division Technical Release 55, Second edition. Washington, D. C.: U.S. Government Printing Office.
- Stanford, J. A. and J. V. Ward. 1993. An ecosystem perspective on alluvial rivers: Connectivity and the hyporheic corridor. *Journal of the North American Benthological Society* 12:48-60.
- Steinman, A.D. Effects of grazers on freshwater benthic algae. Pages 341-373 *in*: R.J. Stevenson, M.L. Bothwell, and R.L. Lowe (editors). *Algal ecology: freshwater benthic ecosystems*. Academic Press.

- Stone, P.A. 2001. Movements and demography of the Sonoran mud turtle, *Kinosternon sonoriense*. *Southwestern Naturalist* 46: 41-53.
- Strahler, A. N. 1957. Quantitative analysis of watershed geomorphology. *Transactions, American Geophysical Union* 38:913-920.
- Strayer, D.L. 1993. Macrohabitats of freshwater mussels (Bivalvia: Unionacea) in streams of the northern Atlantic Slope. *Journal of the North American Benthological Society* 12: 236-246.
- Suberkropp, K., and J.B. Wallace. 1992. Aquatic hyphomycetes in insecticide-treated and untreated streams. *Journal of the North American Benthological Society* 11: 165-171.
- Swanson, F. J., S. V. Gregory, J. R. Sedell, and A. G. Campbell. 1982. Land-water interactions: The riparian zone. pp. 267-291 in: R. L. Edmonds (Editor), *Analysis of coniferous forest ecosystems in the Western U. S.* Hutchinson Ross, Stroudsburg, PA.
- Tarboton, D. G., R. L. Bras, I. Rodrigueziturbe. 1988. The fractal nature of river networks. *Water Resources Research* 24:1317-1323.
- Tasch, P. 1987. *Fossil Conchostraca of the Southern Hemisphere and continental drift.* The Geological Society of America, Boulder, CO, USA.
- Taylor, W.C., and R.J. Hickey. 1992. Habitat, evolution, and speciation in Isoetes. *Annals of the Missouri Botanical Garden* 79: 613-622.
- Tiedje, J.M., S. Brempong, K. Nusslein, T.L. Marsh, and S.J. Flynn. 1999. Opening the black box of soil microbial diversity. *Applied Soil Ecology* 13: 109-122.
- Tiner, R.W., H.C. Bergquist, G.P. DeAlessio, and M.J. Starr. 2002. Geographically isolated wetlands: A preliminary assessment of their characteristics and status in selected areas of the United States. USDA Fish and Wildlife Service, Northeast Region, Hadley, MA. http://wetlands.fws.gov/Pubs_Reports/isolated/report.htm
- Tolland, L. J. G. Cathcart, and S. O. D. Russell. 1998. Estimating the Q100 in British Columbia: A practical problem in forest hydrology. *Journal of the American Water Resources Association* 34:787-794.
- Tramer, E.J. 1977. Catastrophic mortality of stream fishes trapped in shrinking pools. *American Midland Naturalist* 97: 469-478.
- Trimble, S. W. 1997. Contribution of stream channel erosion to sediment yield from an urbanizing watershed. *Science* 278:1442-1444.

- Triska, F. J., and J. R. Sedell. 1976. Decomposition of four species of leaf litter in response to nitrate manipulation. *Ecology* 57:783-792.
- Tsui, C.K.M., K.D. Hyde, and I.J. Hotchkiss. 2001. Longitudinal and temporal distribution of freshwater ascomycetes and dematiaceous hyphomycetes on submerged wood in the Lam Tsuen River, Hong Kong. *Journal of the North American Benthological Society* 20: 533-549.
- United States Fish and Wildlife Service. 1998. Permitted stream losses due to valley filling in Kentucky, Pennsylvania, Virginia, and West Virginia: A partial inventory. Pennsylvania Ecological Services Field Office, State College, PA.
- Vannote R. L., W. G. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences* 37:130-137.
- Vincent, J.B. and J.W. Gibbons. 1995. Terrestrial buffer zones and wetland conservation: A case study of freshwater turtles in a Carolina bay. *Conservation Biology* 9(6): 1365-1369.
- Voelz, N.J., and J.V. McArthur. 2000. An exploration of factors influencing lotic insect species richness. *Biodiversity and Conservation* 9: 1543-1570.
- Wahbe, T.R., and F.L. Bunnell. 2001. Preliminary observations on movements of tailed frog tadpoles (*Ascaphus truei*) in streams through harvested and natural forests. *Northwest Science* 75: 77-83.
- Wallace, J. B., J. R. Webster, and J. L. Meyer. 1995. The influence of log additions on physical and biotic characteristics of a mountain stream. *Canadian Journal of Fisheries and Aquatic Sciences* 52:2120-2137.
- Ward, A. D. and W. J. Elliot. 1995. *Environmental hydrology*. Boca Raton: Lewis Publishers.
- Ward, G. M., and K. W. Cummins. 1979. Effects of food quality on growth of a stream detritivore, *Paratendipes albimanus* (Meigen) (Diptera: Chronomidae). *Ecology* 60:57-64.
- Waterhouse, F.L., A.S. Harestad, and P.K. Ott. 2002. Use of small streams and forest gaps for breeding habitats by winter wrens in coastal British Columbia. *Northwest Science* 76: 335-346.
- Webster, J. R., and J. L. Meyer. 1997a. Stream organic matter budgets. *Journal of the North American Benthological Society* 16:3-161
- Webster, J. R., and J. L. Meyer. 1997b. Organic matter budgets for streams: a synthesis. pages 141-161 in Webster and J. L. Meyer (editors). *Stream organic matter budgets*. *Journal of the North American Benthological Society* 16:3-161.

- Webster, J. R., E. F. Benfield, T. P. Ehrman, M. A. Schaeffer, J. L. Tank, J. J. Hutchens, and D. J. D'Angelo. 1999. What happens to allochthonous material that falls into streams? A synthesis of new and published information from Coweeta. *Freshwater Biology* 41:687-705.
- Weller, C.M., M.C. Watzin, and D. Wang. 1996. The role of wetlands in reducing phosphorus loading to surface water in eight watersheds in the Lake Champlain Basin. *Environmental Management* 20:731-739.
- Wenninger, E.J. and W.F. Fagan. 2000. Effect of river flow manipulation on wolf spider assemblages at three desert riparian sites. *Journal of Arachnology* 28: 115 - 122.
- Whiteside, B.G., and R.M. McNatt. 1972. Fish species diversity in relation to stream order and physicochemical conditions in the Plum Creek drainage basin. *American Midland Naturalist* 88: 90-101.
- Whiting P. J. and J. B. Bradley. 1993. A process-based classification system for headwaters streams. *Earth Surface Processes and Landforms* 18:603-612.
- Whiting, P.J. and M. Pomeranets. 1997. A numerical study of bank storage and its contribution to streamflow. *Journal of Hydrology* 202:121-136.
- Whittaker, R. H., and G. E. Likens. 1973. Carbon in the biota. *Brookhaven Symposium in Biology*. 24:281-302.
- Whittow, J. 1984. *Dictionary of Physical Geography*. New York: Penguin Books.
- Wiebe, K.L., and K. Martin. 1998. Seasonal use by birds of stream-side riparian habitat in coniferous forest of northcentral British Columbia. *Ecography* 21: 124-134.
- Wiest, J.A., Jr. 1982. Anuran succession at temporary ponds in a post oak-savanna region of Texas. Pp. 39-47 *In* N.J. Scott, Jr., ed. *Herpetological communities*. U.S. Fish and Wildlife Service, Washington, D.C.
- Wilbur, H.M. 1987. Regulation of structure in complex systems: Experimental temporary pond communities. *Ecology* 68:1437-1452.
- Wilde, G.R., and K.G. Ostrand. 1999. Changes in the fish assemblage of an intermittent prairie stream upstream from a Texas impoundment. *Texas Journal of Science* 51: 203-210.
- Williams, D.D. 1987. *The ecology of temporary waters*. Croom Helm and Timber Press, London, U. K. , Sydney, AU, and Portland, OR, USA.
- Williams, D.D., N.E. Williams, and Y. Cao. 1997. Spatial differences in macroinvertebrate community structure in springs in southeastern Ontario in relation to their chemical and physical environments. *Canadian Journal of Zoology* 75: 1404-1414.

- Wipfli, M. S., and D. P. Gregovich. 2002. Export of invertebrates and detritus from fishless headwater streams in southeastern Alaska: implications for downstream salmonid production. *Freshwater Biology* 47:957-969.
- Wolman, M. G. and J. P. Miller. 1960. Magnitude and frequency of forces in geomorphic processes. *Journal of Geology* 68:54-74.
- Woltemade, C. J. 1994. Form and process: Fluvial geomorphology and flood-flow interaction, Grant River, Wisconsin. *Annals of the Association of American Geographers* 84:462-479.
- Woltemade, C. J. and K. W. Potter. 1994. A watershed modeling analysis of fluvial geomorphic influences on flood peak attenuation. *Water Resources Research* 30:1933-1942.
- Zedler, J.B. 2003. Wetlands at your service: reducing impacts of agriculture at the watershed scale. *Frontiers in Ecology* 1 (2): 65-72.
- Zedler, P.H. 1987. The ecology of southern California vernal pools: A community profile, Biological Report 85(7.11). U. S. Fish and Wildlife Service, National Wetlands. Research Center, Washington, D. C., USA.
- Zedler, P.H. In press. Vernal pools and the concept of isolated wetlands. *Wetlands*.
- Zedler, P.H., C.K. Frazier, and C. Black. 1993. Habitat creation as a strategy in ecosystem preservation: An example from vernal pools in San Diego County. p. 239-47. *In* J. E. Keeley (Editor) *Interface between ecology and land development in California*. Southern California Academy of Sciences, Los Angeles.
- Zimmerman, A. and M. Church. 2001. Channel morphology, gradient profiles, and bed stresses during flood in a step-pool channel. *Geomorphology* 40:311-327.